

Chapter 10

Innate and Adaptive Immunity in the Skin

Robert L. Modlin, Lloyd S. Miller, Christine Bangert, & Georg Stingl

REFERENCES

1. Metschnikoff E: Ueber eine Sprosspilzkrankheit der Daphnien. Beitrag zur Lehre ueber den Kampf der Phagozyten gegen Krankheitserreger. *Archiv f pathologische Anatomie und Physiologie und f klinische Medizin* **96**:177, 1884
2. Elias PM: Stratum corneum defensive functions: An integrated view. *J Invest Dermatol* **125**:183, 2005
3. Gasque P: Complement: A unique innate immune sensor for danger signals. *Mol Immunol* **41**:1089, 2004
4. Braff MH et al: Cutaneous defense mechanisms by antimicrobial peptides. *J Invest Dermatol* **125**:9, 2005
5. Schaubert J, Gallo RL: Antimicrobial peptides and the skin immune defense system. *J Allergy Clin Immunol* **122**:261, 2008
6. Harder J et al: A peptide antibiotic from human skin. *Nature* **387**:861, 1997
7. Harder J et al: Isolation and characterization of human beta-defensin-3, a novel human inducible peptide antibiotic. *J Biol Chem* **276**:5707, 2001
8. Yang RB et al: Signaling events induced by lipopolysaccharide-activated toll-like receptor 2. *J Immunol* **163**:639, 1999
9. Nizet V et al: Innate antimicrobial peptide protects the skin from invasive bacterial infection. *Nature* **414**:454, 2001
10. Ong PY et al: Endogenous antimicrobial peptides and skin infections in atopic dermatitis. *N Engl J Med* **347**:1151, 2002
11. Lopez-Garcia B et al: Anti-fungal activity of cathelicidins and their potential role in *Candida albicans* skin infection. *J Invest Dermatol* **125**:108, 2005
12. Howell MD et al: Cathelicidin deficiency predisposes to eczema herpeticum. *J Allergy Clin Immunol* **117**:836, 2006
13. Howell MD et al: Cytokine milieu of atopic dermatitis skin subverts the innate immune response to vaccinia virus. *Immunity* **24**:341, 2006
14. De Y et al: LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl peptide receptor-like 1 (FPRL1) as a receptor to chemoattract human peripheral blood neutrophils, monocytes, and T cells. *J Exp Med* **192**:1069, 2000
15. Murakami M et al: Postsecretory processing generates multiple cathelicidins for enhanced topical antimicrobial defense. *J Immunol* **172**:3070, 2004
16. Liu PT et al: Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science* **311**:1770, 2006
17. Schaubert J et al: Injury enhances TLR2 function and antimicrobial peptide expression through a vitamin D-dependent mechanism. *J Clin Invest* **117**:803, 2007
18. Yamasaki K et al: Increased serine protease activity and cathelicidin promotes skin inflammation in rosacea. *Nat Med* **13**:975, 2007
19. Lande R et al: Plasmacytoid dendritic cells sense self-DNA coupled with antimicrobial peptide. *Nature* **449**:564, 2007
20. Glaeser R et al: Antimicrobial psoriasin (S100A7) protects human skin from *Escherichia coli* infection. *Nat Immunol* **6**:57, 2005
21. Michalek M et al: The human antimicrobial protein psoriasin acts by permeabilization of bacterial membranes. *Dev Comp Immunol* **33**:740, 2009
22. Abtin A et al: Flagellin is the principal inducer of the antimicrobial peptide S100A7c (psoriasin) in human epidermal keratinocytes exposed to *Escherichia coli*. *FASEB J* **22**:2168, 2008
23. Jinquan T et al: Psoriasin: A novel chemotactic protein. *J Invest Dermatol* **107**:5, 1996
24. Harder J, Schroeder JM: RNase 7, a novel innate immune defense antimicrobial protein of healthy human skin. *J Biol Chem* **277**:46779, 2002
25. Zanger P et al: Constitutive expression of the antimicrobial peptide RNase 7 is associated with *Staphylococcus aureus* infection of the skin. *J Infect Dis* **200**:1907, 2009

26. Schittek B et al: Dermcidin: A novel human antibiotic peptide secreted by sweat glands. *Nat Immunol* **2**:1133, 2001
27. Senyurek I et al: Dermcidin-derived peptides show a different mode of action than the cathelicidin LL-37 against *Staphylococcus aureus*. *Antimicrob Agents Chemother* **53**:2499, 2009
28. Kalinski P et al: IL-12-deficient dendritic cells, generated in the presence of prostaglandin E₂, promote type 2 cytokine production in maturing human naive T helper cells. *J Immunol* **159**:28, 1997
29. Bickers DR, Athar M: Oxidative stress in the pathogenesis of skin disease. *J Invest Dermatol* **126**:2565, 2006
30. Medzhitov R et al: A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* **388**:394, 1997
31. Lemaitre B et al: The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* **86**:973, 1996
32. Poltorak A et al: Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: Mutations in *Tlr4* gene. *Science* **282**:2085, 1998
33. Klesney-Tait J et al: The TREM receptor family and signal integration. *Nat Immunol* **7**:1266, 2006
34. Crocker PR: Siglecs in innate immunity. *Curr Opin Pharmacol* **5**:431, 2005
35. Robinson MJ et al: Myeloid C-type lectins in innate immunity. *Nat Immunol* **7**:1258, 2006
36. Krutzik SR et al: TLR activation triggers the rapid differentiation of monocytes into macrophages and dendritic cells. *Nat Med* **11**:653, 2005
37. Mohamadzadeh M et al: Activation of triggering receptor expressed on myeloid cells-1 on human neutrophils by marburg and ebola viruses. *J Virol* **80**:7235, 2006
38. Akira S et al: Pathogen recognition and innate immunity. *Cell* **124**:783, 2006
39. Brightbill HD et al: Host defense mechanisms triggered by microbial lipoproteins through toll-like receptors. *Science* **285**:732, 1999
40. Aliprantis AO et al: Cell activation and apoptosis by bacterial lipoproteins through Toll-like receptor-2. *Science* **285**:736, 1999
41. Takeuchi O et al: Discrimination of bacterial lipoproteins by Toll-like receptor 6. *Int Immunol* **13**:933, 2001
42. Takeuchi O et al: Role of TLR1 in mediating immune response to microbial lipoproteins. *J Immunol* **169**:10, 2002
43. Hemmi H et al: A Toll-like receptor that recognizes bacterial DNA. *Nature* **408**:740, 2000
44. Hayashi F et al: The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* **410**:1099, 2001
45. Alexopoulou L et al: Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* **413**:732, 2001
46. Diebold SS et al: Innate antiviral responses by means of TLR7-mediated recognition of single-stranded RNA. *Science* **303**:1529, 2004
47. Staeger H et al: Human toll-like receptors 2 and 4 are targets for deactivation of mononuclear phagocytes by interleukin-4. *Immunol Lett* **71**:1, 2000
48. Medzhitov R, Janeway CA, Jr.: Innate immunity: Impact on the adaptive immune response. *Curr Opin Immunol* **9**:4, 1997
49. Yang RB et al: Toll-like receptor-2 mediates lipopolysaccharide-induced cellular signalling. *Nature* **395**:284, 1998
50. Takeuchi O et al: Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components. *Immunity* **11**:443, 1999
51. Hirschfeld M et al: Cutting edge: Inflammatory signaling by *borrelia burgdorferi* lipoproteins is mediated by toll-like receptor 2. *J Immunol* **163**:2382, 1999
52. Hou L et al: Toll-like receptor 4-deficient mice have reduced bone destruction following mixed anaerobic infection. *Infect Immun* **68**:4681, 2000
53. Gately MK et al: Regulation of human lymphocyte proliferation by a heterodimeric cytokine, IL-12 (cytotoxic lymphocyte maturation factor). *J Immunol* **147**:874, 1991
54. Thoma-Uszynski S et al: Activation of Toll-like receptor 2 on human dendritic cells triggers induction of IL-12 but not IL-10. *J Immunol* **165**:3804, 2000
55. Verreck FA et al: Human IL-23-producing type 1 macrophages promote but IL-10-producing type 2 macrophages subvert immunity to (myco) bacteria. *Proc Natl Acad Sci U S A* **101**:4560, 2004
56. Roses RE et al: Differential production of IL-23 and IL-12 by myeloid-derived dendritic cells in response to TLR agonists. *J Immunol* **181**:5120, 2008

57. Blander JM, Medzhitov R: Regulation of phagosome maturation by signals from toll-like receptors. *Science* **304**:1014, 2004
58. Doyle SE et al: Toll-like receptors induce a phagocytic gene program through p38. *J Exp Med* **199**:81, 2004
59. Krutzik SR et al: IL-15 links TLR2/1-induced macrophage differentiation to the vitamin D-dependent antimicrobial pathway. *J Immunol* **181**:7115, 2008
60. Hertz CJ et al: Microbial lipopeptides stimulate dendritic cell maturation via TLR2. *J Immunol* **166**:2444, 2000
61. Wang TT et al: Cutting edge: 1,25-dihydroxyvitamin D3 is a direct inducer of antimicrobial peptide gene expression. *J Immunol* **173**:2909, 2004
62. Martineau AR et al: IFN-g- and TNF-independent vitamin D-inducible human suppression of mycobacteria: The Role of Cathelicidin LL-37. *J Immunol* **178**:7190, 2007
63. Liu PT et al: Cutting edge: Vitamin D-mediated human antimicrobial activity against *Mycobacterium tuberculosis* is dependent on the induction of cathelicidin. *J Immunol* **179**:2060, 2007
64. Doyle S et al: IRF3 mediates a TLR3/TLR4-specific antiviral gene program. *Immunity* **17**:251, 2002
65. Kawai T et al: Lipopolysaccharide stimulates the MyD88-independent pathway and results in activation of IFN-regulatory factor 3 and the expression of a subset of lipopolysaccharide-inducible genes. *J Immunol* **167**:5887, 2001
66. Kawai T et al: Interferon-alpha induction through Toll-like receptors involves a direct interaction of IRF7 with MyD88 and TRAF6. *Nat Immunol* **5**:1061, 2004
67. Kim J et al: Activation of toll-like receptor 2 in acne triggers inflammatory cytokine responses. *J Immunol* **169**:1535, 2002
68. Aliprantis AO et al: The apoptotic signaling pathway activated by Toll-like receptor-2. *Embo J* **19**:3325, 2000
69. Kriehuber E et al: Balance between NF-kappaB and JNK/AP-1 activity controls dendritic cell life and death. *Blood* **106**:175, 2005
70. Noss EH et al: Toll-like receptor 2-dependent inhibition of macrophage class II MHC expression and antigen processing by 19-kDa lipoprotein of *Mycobacterium tuberculosis*. *J Immunol* **167**:910, 2001
71. Arbour NC et al: TLR4 mutations are associated with endotoxin hyporesponsiveness in humans. *Nat Genet* **25**:187, 2000
72. Strober W et al: Signalling pathways and molecular interactions of NOD1 and NOD2. *Nat Rev Immunol* **6**:9, 2006
73. Fritz JH et al: Nod-like proteins in immunity, inflammation and disease. *Nat Immunol* **7**:1250, 2006
74. Delbridge LM, O'Riordan MX: Innate recognition of intracellular bacteria. *Curr Opin Immunol* **19**:10, 2007
75. Martinon F et al: The inflammasomes: Guardians of the body. *Annu Rev Immunol* **27**:229, 2009
76. Hruz P et al: NOD2 contributes to cutaneous defense against *Staphylococcus aureus* through alpha-toxin-dependent innate immune activation. *Proc Natl Acad Sci U S A* **106**:12873, 2009
77. Hugot JP et al: Association of NOD2 leucine-rich repeat variants with susceptibility to Crohn's disease. *Nature* **411**:599, 2001
78. Miceli-Richard C et al: CARD15 mutations in Blau syndrome. *Nat Genet* **29**:19, 2001
79. Kanazawa N et al: Early-onset sarcoidosis and CARD15 mutations with constitutive nuclear factor-kappaB activation: Common genetic etiology with Blau syndrome. *Blood* **105**:1195, 2005
80. Zhang FR et al: Genomewide association study of leprosy. *N Engl J Med* **361**:2609, 2009
81. Weidinger S et al: Association of NOD1 polymorphisms with atopic eczema and related phenotypes. *J Allergy Clin Immunol* **116**:177, 2005
82. Mosser DM, Edwards JP: Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* **8**:958, 2008
83. Gordon S: Alternative activation of macrophages. *Nat Rev Immunol* **3**:23, 2003
84. Montoya D et al: Divergence of macrophage phagocytic and antimicrobial programs in leprosy. *Cell Host Microbe* **6**:343, 2009
85. van Kooten C, Banchereau J: CD40-CD40 ligand. *J Leukoc Biol* **67**:2, 2000
86. Raulet DH: Interplay of natural killer cells and their receptors with the adaptive immune response. *Nat Immunol* **5**:996, 2004
87. Bauer S et al: Activation of NK cells and T cells by NKG2D, a receptor for stress-inducible MICA. *Science* **285**:727, 1999
88. Groh V et al: Broad tumor-associated expression and recognition by tumor-derived gamma delta T cells of MICA and MICB. *Proc Natl Acad Sci U S A* **96**:6879, 1999

89. Chan CW et al: Interferon-producing killer dendritic cells provide a link between innate and adaptive immunity. *Nat Med* **12**:207, 2006
90. Taieb J et al: A novel dendritic cell subset involved in tumor immunosurveillance. *Nat Med* **12**:214, 2006
91. Kupper TS, Groves RW: The interleukin-1 axis and cutaneous inflammation. *J Invest Dermatol* **105**:62S, 1995
92. Cerwenka A, Swain SL: TGF-beta1: Immunosuppressant and viability factor for T lymphocytes. *Microbes Infect* **1**:1291, 1999
93. Strobl H et al: TGF-beta 1 promotes in vitro development of dendritic cells from CD34 +hemopoietic progenitors. *J Immunol* **157**:1499, 1996
94. Blauvelt A et al: Interleukin-15 mRNA is expressed by human keratinocytes Langerhans cells, and blood-derived dendritic cells and is downregulated by ultraviolet B radiation. *J Invest Dermatol* **106**:1047, 1996
95. Stoll S et al: Production of IL-18 (IFN-gamma-inducing factor) messenger RNA and functional protein by murine keratinocytes. *J Immunol* **159**:298, 1997
96. Enk AH, Katz SI: Identification and induction of keratinocyte-derived IL-10. *J Immunol* **149**:92, 1992
97. Lebre MC et al: Double-stranded RNA-exposed human keratinocytes promote Th1 responses by inducing a Type-1 polarized phenotype in dendritic cells: Role of keratinocyte-derived tumor necrosis factor alpha, type I interferons, and interleukin-18. *J Invest Dermatol* **120**:990, 2003
98. Soumelis V et al: Human epithelial cells trigger dendritic cell mediated allergic inflammation by producing TSLP. *Nat Immunol* **3**:673, 2002
99. Harrington LE et al: Expanding the effector CD4 T-cell repertoire: The Th17 lineage. *Curr Opin Immunol* **18**:349, 2006
100. Enk AH, Katz SI: Early molecular events in the induction phase of contact sensitivity. *Proc Natl Acad Sci U S A* **89**:1398, 1992
101. Kock A et al: Human keratinocytes are a source for tumor necrosis factor alpha: Evidence for synthesis and release upon stimulation with endotoxin or ultraviolet light. *J Exp Med* **172**:1609, 1990
102. Fenjves ES et al: Systemic distribution of apolipoprotein E secreted by grafts of epidermal keratinocytes: Implications for epidermal function and gene therapy. *Proc Natl Acad Sci U S A* **86**:8803, 1989
103. Miller LS et al: TGF-alpha regulates TLR expression and function on epidermal keratinocytes. *J Immunol* **174**:6137, 2005
104. Kollisch G et al: Various members of the Toll-like receptor family contribute to the innate immune response of human epidermal keratinocytes. *Immunology* **114**:531, 2005
105. Lebre MC et al: Human keratinocytes express functional Toll-like receptor 3, 4, 5, and 9. *J Invest Dermatol* **127**:331, 2007
106. Nestle FO et al: Skin immune sentinels in health and disease. *Nat Rev Immunol* **9**:679, 2009
107. Reiss Y et al: CC chemokine receptor (CCR)4 and the CCR10 ligand cutaneous T cell-attracting chemokine (CTACK) in lymphocyte trafficking to inflamed skin. *J Exp Med* **194**:1541, 2001
108. Charbonnier AS et al: Macrophage inflammatory protein 3alpha is involved in the constitutive trafficking of epidermal langerhans cells. *J Exp Med* **190**:1755, 1999
109. Hedrick MN et al: CCR6 is required for IL-23-induced psoriasis-like inflammation in mice. *J Clin Invest* **119**:2317, 2009
110. Harper EG et al: Th17 cytokines stimulate CCL20 expression in keratinocytes in vitro and in vivo: Implications for psoriasis pathogenesis. *J Invest Dermatol* **129**:2175, 2009
111. Liang SC et al: Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* **203**:2271, 2006
112. Peric M et al: IL-17A enhances vitamin D3-induced expression of cathelicidin antimicrobial peptide in human keratinocytes. *J Immunol* **181**:8504, 2008
113. Caruso R et al: Involvement of interleukin-21 in the epidermal hyperplasia of psoriasis. *Nat Med* **15**:1013, 2009
114. Zheng Y et al: Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* **445**:648, 2007
115. Karp CL et al: Mechanism of suppression of cell-mediated immunity by measles virus. *Science* **273**:228, 1996
116. Chen K et al: Immunoglobulin D enhances immune surveillance by activating antimicrobial, proinflammatory and B cell-stimulating programs in basophils. *Nat Immunol* **10**:889, 2009
117. von Boehmer H: Selection of the T-cell repertoire: Receptor-controlled checkpoints in T-cell development. *Adv Immunol* **84**:201, 2004

118. Ferrick DA et al: Differential production of interferon-gamma and interleukin-4 in response to Th1- and Th2-stimulating pathogens by gamma delta T cells in vivo. *Nature* **373**:255, 1995
119. Shibata K et al: Resident Vdelta1+ gammadelta T cells control early infiltration of neutrophils after Escherichia coli infection via IL-17 production. *J Immunol* **178**:4466, 2007
120. Fenoglio D et al: Vdelta1 T lymphocytes producing IFN-gamma and IL-17 are expanded in HIV-1-infected patients and respond to Candida albicans. *Blood* **113**:6611, 2009
121. Peng MY et al: Interleukin 17-producing gamma delta T cells increased in patients with active pulmonary tuberculosis. *Cell Mol Immunol* **5**:203, 2008
122. Ellmeier W et al: The regulation of CD4 and CD8 coreceptor gene expression during T cell development. *Annu Rev Immunol* **17**:523, 1999
123. Takahama Y: Journey through the thymus: Stromal guides for T-cell development and selection. *Nat Rev Immunol* **6**:127, 2006
124. Stenger S et al: Differential effects of cytolytic T cell subsets on intracellular infection. *Science* **276**:1684, 1997
125. Berke G: The binding and lysis of target cells by cytotoxic lymphocytes: Molecular and cellular aspects. *Annu Rev Immunol* **12**:735, 1994
126. Kagi D et al: Cytotoxicity mediated by T cells and natural killer cells is greatly impaired in perforin-deficient mice. *Nature* **369**:31, 1994
127. Stenger S et al: An antimicrobial activity of cytolytic T cells mediated by granulysin. *Science* **282**:121, 1998
128. Palermo B et al: Specific cytotoxic T lymphocyte responses against Melan-A/MART1, tyrosinase and gp100 in vitiligo by the use of major histocompatibility complex/peptide tetramers: The role of cellular immunity in the etiopathogenesis of vitiligo. *J Invest Dermatol* **117**:326, 2001
129. Wang R et al: Interactions between double positive thymocytes and high affinity ligands presented by cortical epithelial cells generate double negative thymocytes with T cell regulatory activity. *Proc Natl Acad Sci U S A* **99**:2181, 2002
130. Kubota H et al: CD3+4-8- alpha beta T cell population with biased T cell receptor V gene usage. Presence in bone marrow and possible involvement of IL-3 for their extrathymic development. *J Immunol* **149**:1143, 1992
131. Strober S et al: Cloned natural suppressor cell lines express the CD3+CD4-CD8- surface phenotype and the alpha, beta heterodimer of the T cell antigen receptor. *J Immunol* **143**:1118, 1989
132. Zhang ZX et al: Identification of a previously unknown antigen-specific regulatory T cell and its mechanism of suppression. *Nat Med* **6**:782, 2000
133. Fischer K et al: Isolation and characterization of human antigen-specific TCR alpha beta +CD4(-) CD8- double-negative regulatory T cells. *Blood* **105**:2828, 2005
134. Ford MS et al: The immune regulatory function of lymphoproliferative double negative T cells in vitro and in vivo. *J Exp Med* **196**:261, 2002
135. Ford McIntyre MS et al: Cutting edge: In vivo trogocytosis as a mechanism of double negative regulatory T cell-mediated antigen-specific suppression. *J Immunol* **181**:2271, 2008
136. Young KJ et al: Antitumor activity mediated by double-negative T cells. *Cancer Res* **63**:8014, 2003
137. Martinez C et al: Functional double-negative T cells in the periphery express T cell receptor V beta gene products that cause deletion of single-positive T cells. *Eur J Immunol* **23**:250, 1993
138. Nascimbeni M et al: Peripheral CD4(+)CD8(+) T cells are differentiated effector memory cells with antiviral functions. *Blood* **104**:478, 2004
139. Jimenez E et al: Rat peripheral CD4+CD8+ T lymphocytes are partially immunocompetent thymus-derived cells that undergo post-thymic maturation to become functionally mature CD4+ T lymphocytes. *J Immunol* **168**:5005, 2002
140. Weiss L et al: Persistent expansion, in a human immunodeficiency virus-infected person, of V beta-restricted CD4+CD8+ T lymphocytes that express cytotoxicity-associated molecules and are committed to produce interferon-gamma and tumor necrosis factor-alpha. *J Infect Dis* **178**:1158, 1998
141. Sala P et al: Persistent expansions of CD4+ CD8+ peripheral blood T cells. *Blood* **82**:1546, 1993
142. Ober BT et al: Vaccine-induced, pseudorabies virus-specific, extrathymic CD4+CD8+ memory T-helper cells in swine. *J Virol* **72**:4866, 1998
143. Desfrancois J et al: Increased frequency of non-conventional double positive CD4CD8 alphabeta T cells in human breast pleural effusions. *Int J Cancer* **125**:374, 2009
144. Desfrancois J et al: Double positive CD4CD8 alphabeta T cells: A new tumor-reactive population in human melanomas. *PLoS One* **5**:e8437, 2010
145. Krogsgaard M et al: A role for "self" in T-cell activation. *Semin Immunol* **19**:236, 2007

146. Link A et al: Fibroblastic reticular cells in lymph nodes regulate the homeostasis of naive T cells. *Nat Immunol* **8**:1255, 2007
147. Kerdiles YM et al: Foxo1 links homing and survival of naive T cells by regulating L-selectin, CCR7 and interleukin 7 receptor. *Nat Immunol* **10**:176, 2009
148. Sprent J, Surh CD: T cell memory. *Annu Rev Immunol* **20**:551, 2002
149. Lanzavecchia A, Sallusto F: Progressive differentiation and selection of the fittest in the immune response. *Nat Rev Immunol* **2**:982, 2002
150. Tough DF, Sprent J: Turnover of naive- and memory-phenotype T cells. *J Exp Med* **179**:1127, 1994
151. Surh CD, Sprent J: Homeostasis of naive and memory T cells. *Immunity* **29**:848, 2008
152. Sallusto F et al: Central memory and effector memory T cell subsets: Function, generation, and maintenance. *Annu Rev Immunol* **22**:745, 2004
153. Wherry EJ et al: Lineage relationship and protective immunity of memory CD8 T cell subsets. *Nat Immunol* **4**:225, 2003
154. Mitchell DM et al: Distinct roles for IL-2 and IL-15 in the differentiation and survival of CD8⁺ effector and memory T cells. *J Immunol* **184**:6719, 2010
155. Sallusto F, Lanzavecchia A: Heterogeneity of CD4⁺ memory T cells: Functional modules for tailored immunity. *Eur J Immunol* **39**:2076, 2009
156. Duhon T et al: Production of interleukin 22 but not interleukin 17 by a subset of human skin-homing memory T cells. *Nat Immunol* **10**:857, 2009
157. Veldhoen M et al: Transforming growth factor-beta 'reprograms' the differentiation of T helper 2 cells and promotes an interleukin 9-producing subset. *Nat Immunol* **9**:1341, 2008
158. Eyerich S et al: Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* **119**:3573, 2009
159. Lu LF et al: Mast cells are essential intermediaries in regulatory T-cell tolerance. *Nature* **442**:997, 2006
160. Mosmann TR et al: Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* **136**:2348, 1986
161. Abbas AK et al: Functional diversity of helper T lymphocytes. *Nature* **383**:787, 1996
162. Szabo SJ et al: Distinct effects of T-bet in TH1 lineage commitment and IFN-gamma production in CD4 and CD8 T cells. *Science* **295**:338, 2002
163. Zheng W, Flavell RA: The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* **89**:587, 1997
164. Lee GR et al: Regulation of IL-4 gene expression by distal regulatory elements and GATA-3 at the chromatin level. *Immunity* **14**:447, 2001
165. Heinzel FP et al: Production of interferon gamma, interleukin 2, interleukin 4, and interleukin 10 by CD4⁺ lymphocytes in vivo during healing and progressive murine leishmaniasis. *Proc Natl Acad Sci USA* **88**:7011, 1991
166. Pearce EJ et al: Downregulation of Th1 cytokine production accompanies induction of Th2 responses by a parasitic helminth, *Schistosoma mansoni*. *J Exp Med* **173**:159, 1991
167. Pond L et al: Evidence for differential induction of helper T cell subsets during *Trichinella spiralis* infection. *J Immunol* **143**:4232, 1989
168. Gately MK et al: The interleukin-12/interleukin-12-receptor system: Role in normal and pathologic immune responses. *Annu Rev Immunol* **16**:495, 1998
169. Korn T et al: IL-17 and Th17 cells. *Annu Rev Immunol* **27**:485, 2009
170. Ivanov, II et al: The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17⁺ T helper cells. *Cell* **126**:1121, 2006
171. Kleiweietfeld M et al: CCR6 expression defines regulatory effector/memory-like cells within the CD25⁺CD4⁺ T-cell subset. *Blood* **105**:2877, 2005
172. Cosmi L et al: Human interleukin 17-producing cells originate from a CD161⁺CD4⁺ T cell precursor. *J Exp Med* **205**:1903, 2008
173. Wilson NJ et al: Development, cytokine profile and function of human interleukin 17-producing helper T cells. *Nat Immunol* **8**:950, 2007
174. Manel N et al: The differentiation of human T(H)-17 cells requires transforming growth factor-beta and induction of the nuclear receptor ROR γ . *Nat Immunol* **9**:641, 2008
175. Kleinschek MA et al: Circulating and gut-resident human Th17 cells express CD161 and promote intestinal inflammation. *J Exp Med* **206**:525, 2009
176. Zou W, Restifo NP: T(H)17 cells in tumour immunity and immunotherapy. *Nat Rev Immunol* **10**:248, 2010
177. He D et al: CD8⁺IL-17-producing T cells are important in effector functions for the elicitation of contact hypersensitivity responses. *J Immunol* **177**:6852, 2006

178. Boniface K et al: IL-22 inhibits epidermal differentiation and induces proinflammatory gene expression and migration of human keratinocytes. *J Immunol* **174**:3695, 2005
179. Bendelac A et al: The biology of NKT cells. *Annu Rev Immunol* **25**:297, 2007
180. Jiang H, Chess L: Regulation of immune responses by T cells. *N Engl J Med* **354**:1166, 2006
181. Shevach EM et al: The lifestyle of naturally occurring CD4⁺ CD25⁺ Foxp3⁺ regulatory T cells. *Immunol Rev* **212**:60, 2006
182. Shevach EM: Mechanisms of foxp3⁺ T regulatory cell-mediated suppression. *Immunity* **30**:636, 2009
183. Voo KS et al: Identification of IL-17-producing FOXP3⁺ regulatory T cells in humans. *Proc Natl Acad Sci U S A* **106**:4793, 2009
184. Kuhl AA et al: Human peripheral gammadelta T cells possess regulatory potential. *Immunology* **128**:580, 2009
185. Breitfeld D et al: Follicular B helper T cells express CXC chemokine receptor 5, localize to B cell follicles, and support immunoglobulin production. *J Exp Med* **192**:1545, 2000
186. Schaerli P et al: CXC chemokine receptor 5 expression defines follicular homing T cells with B cell helper function. *J Exp Med* **192**:1553, 2000
187. Fazilleau N et al: Follicular helper T cells: Lineage and location. *Immunity* **30**:324, 2009
188. Campbell DJ, Butcher EC: Rapid acquisition of tissue-specific homing phenotypes by CD4⁽⁺⁾ T cells activated in cutaneous or mucosal lymphoid tissues. *J Exp Med* **195**:135, 2002
189. Sallusto F, Mackay CR: Chemoattractants and their receptors in homeostasis and inflammation. *Curr Opin Immunol* **16**:724, 2004
190. Wakim LM et al: Cutting edge: Local recall responses by memory T cells newly recruited to peripheral nonlymphoid tissues. *J Immunol* **181**:5837, 2008
191. Wakim LM et al: Dendritic cell-induced memory T cell activation in nonlymphoid tissues. *Science* **319**:198, 2008
192. Zhu J et al: Virus-specific CD8⁺ T cells accumulate near sensory nerve endings in genital skin during subclinical HSV-2 reactivation. *J Exp Med* **204**:595, 2007
193. Teraki Y, Shiohara T: IFN-gamma-producing effector CD8⁺ T cells and IL-10-producing regulatory CD4⁺ T cells in fixed drug eruption. *J Allergy Clin Immunol* **112**:609, 2003
194. Clark RA et al: The vast majority of CLA⁺ T cells are resident in normal skin. *J Immunol* **176**:4431, 2006
195. Edele F et al: Cutting edge: Instructive role of peripheral tissue cells in the imprinting of T cell homing receptor patterns. *J Immunol* **181**:3745, 2008
196. Fuhlbrigge RC et al: Cutaneous lymphocyte antigen is a specialized form of PSGL-1 expressed on skin-homing T cells. *Nature* **389**:978, 1997
197. Robert C et al: Interaction of dendritic cells with skin endothelium: A new perspective on immunosurveillance. *J Exp Med* **189**:627, 1999
198. Morales J et al: CTACK, a skin-associated chemokine that preferentially attracts skin-homing memory T cells. *Proc Natl Acad Sci U S A* **96**:14470, 1999
199. Lonsdorf AS et al: Chemokine receptors in T-cell-mediated diseases of the skin. *J Invest Dermatol* **129**:2552, 2009
200. Singh SP et al: Human T cells that are able to produce IL-17 express the chemokine receptor CCR6. *J Immunol* **180**:214, 2008
201. Modlin RL et al: T lymphocyte subsets in the skin lesions of patients with leprosy. *J Am Acad Dermatol* **8**:182, 1983
202. Wood GS et al: Detection of clonal T-cell receptor gamma gene rearrangements in early mycosis fungoides/Sezary syndrome by polymerase chain reaction and denaturing gradient gel electrophoresis (PCR/DGGE). *J Invest Dermatol* **103**:34, 1994
203. Winter D et al: Definition of TCR epitopes for CTL-mediated attack of cutaneous T cell lymphoma. *J Immunol* **171**:2714, 2003
204. Wang XH et al: Selection of T lymphocytes bearing limited T-cell receptor beta chains in the response to a human pathogen. *Proc Natl Acad Sci U S A* **90**:188, 1993
205. Diluvio L et al: Identical TCR beta-chain rearrangements in streptococcal angina and skin lesions of patients with psoriasis vulgaris. *J Immunol* **176**:7104, 2006
206. McCluskey RT et al: Studies on the specificity of the cellular infiltrate in delayed hypersensitivity reactions. *J Immunol* **90**:466, 1963
207. Modlin RL et al: Learning from lesions: Patterns of tissue inflammation in leprosy. *Proc Natl Acad Sci U S A* **85**:1213, 1988
208. Clark RA et al: A novel method for the isolation of skin resident T cells from normal and diseased human skin. *J Invest Dermatol* **126**:1059, 2006

209. Van Voorhis WC et al: Primary and secondary syphilis lesions contain mRNA for Th1 cytokines. *J Infect Dis* **173**:491, 1996
210. Stary G et al: Host defense mechanisms in secondary syphilitic lesions: A role of IFN-gamma/IL-17-producing CD8+ T-cells? *Am J Pathol*, in press 2010
211. Yssel H et al: *Borrelia burgdorferi* activates a T helper type 1-like T cell subset in Lyme arthritis. *J Exp Med* **174**:593, 1991
212. Burchill MA et al: Inhibition of interleukin-17 prevents the development of arthritis in vaccinated mice challenged with *Borrelia burgdorferi*. *Infect Immun* **71**:3437, 2003
213. Kim EJ et al: Immunopathogenesis and therapy of cutaneous T cell lymphoma. *J Clin Invest* **115**:798, 2005
214. Kaporis HG et al: Human basal cell carcinoma is associated with Foxp3+ T cells in a Th2 dominant microenvironment. *J Invest Dermatol* **127**:2391, 2007
215. Ghoreishi M et al: Type 1 interferon signature in the scalp lesions of alopecia areata. *Br J Dermatol* **163**:57, 2010
216. Watts C: Capture and processing of exogenous antigens for presentation on MHC molecules. *Annu Rev Immunol* **15**:821, 1997
217. Pamer E, Cresswell P: Mechanisms of MHC class I-Restricted antigen processing. *Annu Rev Immunol* **16**:323, 1998
218. Hammer GE et al: The final touches make perfect the peptide-MHC class I repertoire. *Immunity* **26**:397, 2007
219. Huang AY et al: Role of bone marrow-derived cells in presenting MHC class I-restricted tumor antigens. *Science* **264**:961, 1994
220. Matsuo M et al: IFN-gamma enables cross-presentation of exogenous protein antigen in human Langerhans cells by potentiating maturation. *Proc Natl Acad Sci U S A* **101**:14467, 2004
221. Stoitzner P et al: Langerhans cells cross-present antigen derived from skin. *Proc Natl Acad Sci U S A* **103**:7783, 2006
222. Gelin C et al: Regulation of MHC II and CD1 antigen presentation: From ubiquity to security. *J Leukoc Biol* **85**:215, 2009
223. Fiebiger E et al: Cytokines regulate proteolysis in major histocompatibility complex class II-dependent antigen presentation by dendritic cells. *J Exp Med* **193**:881, 2001
224. Trombetta ES et al: Activation of lysosomal function during dendritic cell maturation. *Science* **299**:1400, 2003
225. Porcelli SA, Modlin RL: The CD1 system: Antigen-presenting molecules for T cell recognition of lipids and glycolipids. *Annu Rev Immunol* **17**:297, 1999
226. Barral DC, Brenner MB: CD1 antigen presentation: How it works. *Nat Rev Immunol* **7**:929, 2007
227. Steinman RM, Cohn ZA: Identification of a novel cell type in peripheral lymphoid organs of mice. I. Morphology, quantitation, tissue distribution. *J Exp Med* **137**:1142, 1973
228. Stingl G et al: Epidermal Langerhans cells bear Fc and C3 receptors. *Nature* **268**:245, 1977
229. Rowden G et al: Ia antigen expression on human epidermal Langerhans cells. *Nature* **268**:247, 1977
230. Klareskog L et al: Epidermal Langerhans cells express Ia antigens. *Nature* **268**:248, 1977
231. Fogg DK et al: A clonogenic bone marrow progenitor specific for macrophages and dendritic cells. *Science* **311**:83, 2006
232. Naik SH et al: Development of plasmacytoid and conventional dendritic cell subtypes from single precursor cells derived in vitro and in vivo. *Nat Immunol* **8**:1217, 2007
233. Liu K et al: Origin of dendritic cells in peripheral lymphoid organs of mice. *Nat Immunol* **8**:578, 2007
234. Massberg S et al: Immunosurveillance by hematopoietic progenitor cells trafficking through blood, lymph, and peripheral tissues. *Cell* **131**:994, 2007
235. Onai N et al: Identification of clonogenic common Flt3+M-CSFR+ plasmacytoid and conventional dendritic cell progenitors in mouse bone marrow. *Nat Immunol* **8**:1207, 2007
236. Geissmann F et al: Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity* **19**:71, 2003
237. Stumbles PA et al: Regulation of dendritic cell recruitment into resting and inflamed airway epithelium: Use of alternative chemokine receptors as a function of inducing stimulus. *J Immunol* **167**:228, 2001
238. Merad M et al: Langerhans cells renew in the skin throughout life under steady-state conditions. *Nat Immunol* **3**:1135, 2002
239. Merad M et al: Depletion of host Langerhans cells before transplantation of donor alloreactive T cells prevents skin graft-versus-host disease. *Nat Med* **10**:510, 2004

240. Ward BR et al: Local thermal injury elicits immediate dynamic behavioural responses by corneal Langerhans cells. *Immunology* **120**:556, 2007
241. Forster R et al: CCR7 coordinates the primary immune response by establishing functional microenvironments in secondary lymphoid organs. *Cell* **99**:23, 1999
242. Ohl L et al: CCR7 governs skin dendritic cell migration under inflammatory and steady-state conditions. *Immunity* **21**:279, 2004
243. Alvarez D et al: Mechanisms and consequences of dendritic cell migration. *Immunity* **29**:325, 2008
244. Brand CU et al: Studies on human skin lymph containing Langerhans cells from sodium lauryl sulphate contact dermatitis. *J Invest Dermatol* **99**:109S, 1992
245. Tomura M et al: Monitoring cellular movement in vivo with photoconvertible fluorescence protein "Kaede" transgenic mice. *Proc Natl Acad Sci US A* **105**:10871, 2008
246. Ratzinger G et al: Matrix metalloproteinases 9 and 2 are necessary for the migration of Langerhans cells and dermal dendritic cells from human and murine skin. *J Immunol* **168**:4361, 2002
247. Ichiyasu H et al: Matrix metalloproteinase-9-deficient dendritic cells have impaired migration through tracheal epithelial tight junctions. *Am J Respir Cell Mol Biol* **30**:761, 2004
248. Yen JH et al: PGE2-induced metalloproteinase-9 is essential for dendritic cell migration. *Blood* **111**:260, 2008
249. Darmanin S et al: All-trans retinoic acid enhances murine dendritic cell migration to draining lymph nodes via the balance of matrix metalloproteinases and their inhibitors. *J Immunol* **179**:4616, 2007
250. Hugues S et al: Distinct T cell dynamics in lymph nodes during the induction of tolerance and immunity. *Nat Immunol* **5**:1235, 2004
251. Lindquist RL et al: Visualizing dendritic cell networks in vivo. *Nat Immunol* **5**:1243, 2004
252. Mueller DL et al: Clonal expansion versus functional clonal inactivation: A costimulatory signaling pathway determines the outcome of T cell antigen receptor occupancy. *Annu Rev Immunol* **7**:445, 1989
253. Matzinger P: An innate sense of danger. *Ann N Y Acad Sci* **961**:341, 2002
254. Hirano N et al: Engagement of CD83 ligand induces prolonged expansion of CD8+ T cells and preferential enrichment for antigen specificity. *Blood* **107**:1528, 2006
255. Prechtel AT, Steinkasserer A: CD83: An update on functions and prospects of the maturation marker of dendritic cells. *Arch Dermatol Res* **299**:59, 2007
256. Jonuleit H, E. Schmitt: Dendritic cells as a tool to induce anergic and regulatory T cells. *Trends Immunol* **22**:394, 2001
257. Mahnke K et al: Regulatory conversation between antigen presenting cells and regulatory T cells enhance immune suppression. *Cell Immunol* **250**:1, 2007
258. Steinman RM et al: Dendritic cell function in vivo during the steady state: A role in peripheral tolerance. *Ann N Y Acad Sci* **987**:15, 2003
259. Hawiger D et al: Dendritic cells induce peripheral T cell unresponsiveness under steady state conditions in vivo. *J Exp Med* **194**:769, 2001
260. Bonifaz L et al: Efficient targeting of protein antigen to the dendritic cell receptor DEC-205 in the steady state leads to antigen presentation on major histocompatibility complex class I products and peripheral CD8+ T cell tolerance. *J Exp Med* **196**:1627, 2002
261. Probst HC et al: Inducible transgenic mice reveal resting dendritic cells as potent inducers of CD8+ T cell tolerance. *Immunity* **18**:713, 2003
262. Bruder D et al: On the edge of autoimmunity: T-cell stimulation by steady-state dendritic cells prevents autoimmune diabetes. *Diabetes* **54**:3395, 2005
263. Mommaas AM et al: Human epidermal Langerhans cells lack functional mannose receptors and a fully developed endosomal/lysosomal compartment for loading of HLA class II molecules. *Eur J Immunol* **29**:571, 1999
264. Lutz MB, Schuler G: Immature, semi-mature and fully mature dendritic cells: Which signals induce tolerance or immunity? *Trends Immunol* **23**:445, 2002
265. Freeman GJ et al: Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation. *J Exp Med* **192**:1027, 2000
266. Latchman Y et al: PD-L2 is a second ligand for PD-1 and inhibits T cell activation. *Nat Immunol* **2**:261, 2001
267. Tseng SY et al: B7-DC, a new dendritic cell molecule with potent costimulatory properties for T cells. *J Exp Med* **193**:839, 2001
268. Muller G et al: Interleukin-10-treated dendritic cells modulate immune responses of naive and sensitized T cells in vivo. *J Invest Dermatol* **119**:836, 2002

269. Schwarz A et al: Ultraviolet radiation-induced regulatory T cells not only inhibit the induction but can suppress the effector phase of contact hypersensitivity. *J Immunol* **172**:1036, 2004
270. von Bubnoff D et al: Human epidermal langerhans cells express the immunoregulatory enzyme indoleamine 2,3-dioxygenase. *J Invest Dermatol* **123**:298, 2004
271. Hosoi J et al: Regulation of Langerhans cell function by nerves containing calcitonin gene-related peptide. *Nature* **363**:159, 1993
272. Kodali S et al: Vasoactive intestinal peptide modulates Langerhans cell immune function. *J Immunol* **173**:6082, 2004
273. Fukunaga A et al: Src homology 2 domain-containing protein tyrosine phosphatase substrate 1 regulates the induction of Langerhans cell maturation. *Eur J Immunol* **36**:3216, 2006
274. Seiffert K, Granstein RD: Neuroendocrine regulation of skin dendritic cells. *Ann N Y Acad Sci* **1088**:195, 2006
275. Mayerova D et al: Langerhans cells activate naive self-antigen-specific CD8 T cells in the steady state. *Immunity* **21**:391, 2004
276. Kissenpfennig A et al: Dynamics and function of Langerhans cells in vivo: Dermal dendritic cells colonize lymph node areas distinct from slower migrating Langerhans cells. *Immunity* **22**:643, 2005
277. Kaplan DH et al: Epidermal langerhans cell-deficient mice develop enhanced contact hypersensitivity. *Immunity* **23**:611, 2005
278. Cerio R et al: Characterization of factor XIIIa positive dermal dendritic cells in normal and inflamed skin. *Br J Dermatol* **121**:421, 1989
279. Banchereau J, Steinman RM: Dendritic cells and the control of immunity. *Nature* **392**:245, 1998
280. Bursch LS et al: Identification of a novel population of Langerin+ dendritic cells. *J Exp Med* **204**:3147, 2007
281. Ginhoux F et al: Blood-derived dermal langerin+ dendritic cells survey the skin in the steady state. *J Exp Med* **204**:3133, 2007
282. Poulin LF et al: The dermis contains langerin+ dendritic cells that develop and function independently of epidermal Langerhans cells. *J Exp Med* **204**:3119, 2007
283. Romani N et al: Langerhans cells and more: Langerin-expressing dendritic cell subsets in the skin. *Immunol Rev* **234**:120, 2010
284. Langerhans P: Über die Nerven der menschlichen Haut. *Virchows Arch* **44**:325, 1868
285. Kissenpfennig A et al: Disruption of the langerin/CD207 gene abolishes Birbeck granules without a marked loss of Langerhans cell function. *Mol Cell Biol* **25**:88, 2005
286. Schuster C et al: HLA-DR+ leukocytes acquire CD1 antigens in embryonic and fetal human skin and contain functional antigen-presenting cells. *J Exp Med* **206**:169, 2009
287. Tang A et al: Adhesion of epidermal Langerhans cells to keratinocytes mediated by E-cadherin. *Nature* **361**:82, 1993
288. Konrad K, Honigsmann H: Electron microscopic demonstration of a mitotic Langerhans cell in the normal human epidermis. *Arch Dermatol Forsch* **246**:70, 1973
289. Vishwanath M et al: Development of intravital intermittent confocal imaging system for studying langerhans cell turnover. *J Invest Dermatol* **126**:2452, 2006
290. Katz SI et al: Epidermal Langerhans cells are derived from cells originating in bone marrow. *Nature* **282**:324, 1979
291. Larregina AT et al: Dermal-resident CD14+ cells differentiate into Langerhans cells. *Nat Immunol* **2**:1151, 2001
292. Schaeferli P et al: Cutaneous CXCL14 targets blood precursors to epidermal niches for Langerhans cell differentiation. *Immunity* **23**:331, 2005
293. Ginhoux F et al: Langerhans cells arise from monocytes in vivo. *Nat Immunol* **7**:265, 2006
294. Mende I et al: Flk2 +myeloid progenitors are the main source of Langerhans cells. *Blood* **107**:1383, 2006
295. Stingl G et al: Origin and function of epidermal Langerhans cells. *Immunol Rev* **53**:149, 1980
296. Inaba K et al: Immunologic properties of purified epidermal Langerhans cells. Distinct requirements for stimulation of unprimed and sensitized T lymphocytes. *J Exp Med* **164**:605, 1986
297. Hauser C, Katz SI: Activation and expansion of hapten- and protein-specific T helper cells from nonsensitized mice. *Proc Natl Acad Sci U S A* **85**:5625, 1988
298. Romani N et al: Presentation of exogenous protein antigens by dendritic cells to T cell clones. Intact protein is presented best by immature, epidermal Langerhans cells. *J Exp Med* **169**:1169, 1989
299. Kubo A et al: External antigen uptake by Langerhans cells with reorganization of epidermal tight junction barriers. *J Exp Med* **206**:2937, 2009

300. Larsen CP et al: Migration and maturation of Langerhans cells in skin transplants and explants. *J Exp Med* **172**:1483, 1990
301. Schuler G, Steinman RM: Murine epidermal Langerhans cells mature into potent immunostimulatory dendritic cells in vitro. *J Exp Med* **161**:526, 1985
302. Nishibu A et al: Behavioral responses of epidermal Langerhans cells in situ to local pathological stimuli. *J Invest Dermatol* **126**:787, 2006
303. Cumberbatch M et al: Impaired Langerhans cell migration in psoriasis. *J Exp Med* **203**:953, 2006
304. Weiss JM et al: Osteopontin is involved in the initiation of cutaneous contact hypersensitivity by inducing Langerhans and dendritic cell migration to lymph nodes. *J Exp Med* **194**:1219, 2001
305. Kriehuber E et al: Isolation and characterization of dermal lymphatic and blood endothelial cells reveal stable and functionally specialized cell lineages. *J Exp Med* **194**:797, 2001
306. Eberhard Y et al: Up-regulation of the chemokine CCL21 in the skin of subjects exposed to irritants. *BMC Immunol* **5**:7, 2004
307. Meunier L et al: Heterogeneous populations of class II MHC+ cells in human dermal cell suspensions. Identification of a small subset responsible for potent dermal antigen-presenting cell activity with features analogous to Langerhans cells. *J Immunol* **151**:4067, 1993
308. Nestle FO et al: Characterization of dermal dendritic cells obtained from normal human skin reveals phenotypic and functionally distinctive subsets. *J Immunol* **151**:6535, 1993
309. Zaba LC et al: Resident and "inflammatory" dendritic cells in human skin. *J Invest Dermatol* **129**:302, 2009
310. Ochoa MT et al: "Dermal dendritic cells" comprise two distinct populations: CD1+ dendritic cells and CD209+ macrophages. *J Invest Dermatol* **128**:2225, 2008
311. Torocsik D et al: Identification of factor XIII-A as a marker of alternative macrophage activation. *Cell Mol Life Sci* **62**:2132, 2005
312. Zaba LC et al: Normal human dermis contains distinct populations of CD11c+BDCA-1+ dendritic cells and CD163+FXIIIa+ macrophages. *J Clin Invest* **117**:2517, 2007
313. Jongbloed SL et al: Human CD141+ (BDCA-3)+ dendritic cells (DCs) represent a unique myeloid DC subset that cross-presents necrotic cell antigens. *J Exp Med* **207**:1247, 2010
314. Bogunovic M et al: Identification of a radio-resistant and cycling dermal dendritic cell population in mice and men. *J Exp Med* **203**:2627, 2006
315. Angel CE et al: Cutting edge: CD1a+ antigen-presenting cells in human dermis respond rapidly to CCR7 ligands. *J Immunol* **176**:5730, 2006
316. Bennett CL et al: Inducible ablation of mouse Langerhans cells diminishes but fails to abrogate contact hypersensitivity. *J Cell Biol* **169**:569, 2005
317. Streilein JW et al: Depletion of epidermal langerhans cells and Ia immunogenicity from tape-stripped mouse skin. *J Exp Med* **155**:863, 1982
318. Obhrai JS et al: Langerhans cells are not required for efficient skin graft rejection. *J Invest Dermatol* **128**:1950, 2008
319. Schwarz A et al: Langerhans cells are required for UVR-induced immunosuppression. *J Invest Dermatol* **130**:1419, 2010
320. Randolph GJ et al: The CD16(+)(FcγRIII(+)) subset of human monocytes preferentially becomes migratory dendritic cells in a model tissue setting. *J Exp Med* **196**:517, 2002
321. Piccioli D et al: Functional specialization of human circulating CD16 and CD1c myeloid dendritic-cell subsets. *Blood* **109**:5371, 2007
322. Lowes MA et al: Increase in TNF-alpha and inducible nitric oxide synthase-expressing dendritic cells in psoriasis and reduction with efalizumab (anti-CD11a). *Proc Natl Acad Sci USA* **102**:19057, 2005
323. Guttman-Yassky E et al: Major differences in inflammatory dendritic cells and their products distinguish atopic dermatitis from psoriasis. *J Allergy Clin Immunol* **119**:1210, 2007
324. Stary G et al: Tumoricidal activity of TLR7/8-activated inflammatory dendritic cells. *J Exp Med* **204**:1441, 2007
325. Zaba LC et al: Psoriasis is characterized by accumulation of immunostimulatory and Th1/Th17 cell-polarizing myeloid dendritic cells. *J Invest Dermatol* **129**:79, 2009
326. Zaba LC et al: Amelioration of epidermal hyperplasia by TNF inhibition is associated with reduced Th17 responses. *J Exp Med* **204**:3183, 2007
327. Serbina NV et al: TNF/iNOS-producing dendritic cells mediate innate immune defense against bacterial infection. *Immunity* **19**:59, 2003
328. Tezuka H et al: Regulation of IgA production by naturally occurring TNF/iNOS-producing dendritic cells. *Nature* **448**:929, 2007

329. Dockrell DH, Kinghorn GR: Imiquimod and resiquimod as novel immunomodulators. *J Antimicrob Chemother* **48**:751, 2001
330. Palamara F et al: Identification and characterization of pDC-like cells in normal mouse skin and melanomas treated with imiquimod. *J Immunol* **173**:3051, 2004
331. Haider AS et al: Identification of cellular pathways of "type 1," Th17 T cells, and TNF- and inducible nitric oxide synthase-producing dendritic cells in autoimmune inflammation through pharmacogenomic study of cyclosporine A in psoriasis. *J Immunol* **180**:1913, 2008
332. Zaba LC et al: Identification of TNF-related apoptosis-inducing ligand and other molecules that distinguish inflammatory from resident dendritic cells in patients with psoriasis. *J Allergy Clin Immunol* **125**:1261, 2010
333. Wollenberg A et al: Immunomorphological and ultrastructural characterization of Langerhans cells and a novel, inflammatory dendritic epidermal cell (IDEC) population in lesional skin of atopic eczema. *J Invest Dermatol* **106**:446, 1996
334. Wollenberg A et al: Expression and function of the mannose receptor CD206 on epidermal dendritic cells in inflammatory skin diseases. *J Invest Dermatol* **118**:327, 2002
335. Schuller E et al: In situ expression of the costimulatory molecules CD80 and CD86 on langerhans cells and inflammatory dendritic epidermal cells (IDEC) in atopic dermatitis. *Arch Dermatol Res* **293**:448, 2001
336. Novak N et al: FcepsilonRI engagement of Langerhans cell-like dendritic cells and inflammatory dendritic epidermal cell-like dendritic cells induces chemotactic signals and different T-cell phenotypes in vitro. *J Allergy Clin Immunol* **113**:949, 2004
337. Lande R, Gilliet M: Plasmacytoid dendritic cells: Key players in the initiation and regulation of immune responses. *Ann N Y Acad Sci* **1183**:89, 2010
338. Kohrgruber N et al: Survival, maturation, and function of CD11c- and CD11c+ peripheral blood dendritic cells are differentially regulated by cytokines. *J Immunol* **163**:3250, 1999
339. Siegal FP et al: The nature of the principal type 1 interferon-producing cells in human blood. *Science* **284**:1835, 1999
340. Cisse B et al: Transcription factor E2-2 is an essential and specific regulator of plasmacytoid dendritic cell development. *Cell* **135**:37, 2008
341. Gerlini G et al: Massive recruitment of type I interferon producing plasmacytoid dendritic cells in varicella skin lesions. *J Invest Dermatol* **126**:507, 2006
342. Farkas L et al: Plasmacytoid dendritic cells (natural interferon- alpha/beta-producing cells) accumulate in cutaneous lupus erythematosus lesions. *Am J Pathol* **159**:237, 2001
343. Wollenberg A et al: Plasmacytoid dendritic cells: A new cutaneous dendritic cell subset with distinct role in inflammatory skin diseases. *J Invest Dermatol* **119**:1096, 2002
344. Bangert C et al: Immunopathologic features of allergic contact dermatitis in humans: Participation of plasmacytoid dendritic cells in the pathogenesis of the disease? *J Invest Dermatol* **121**:1409, 2003
345. Vanbervliet B et al: The inducible CXCR3 ligands control plasmacytoid dendritic cell responsiveness to the constitutive chemokine stromal cell-derived factor 1 (SDF-1)/CXCL12. *J Exp Med* **198**:823, 2003
346. Kohrgruber N et al: Plasmacytoid dendritic cell recruitment by immobilized CXCR3 ligands. *J Immunol* **173**:6592, 2004
347. Vermi W et al: Role of ChemR23 in directing the migration of myeloid and plasmacytoid dendritic cells to lymphoid organs and inflamed skin. *J Exp Med* **201**:509, 2005
348. Albanesi C et al: Chemerin expression marks early psoriatic skin lesions and correlates with plasmacytoid dendritic cell recruitment. *J Exp Med* **206**:249, 2009
349. Kadowaki N et al: Natural interferon alpha/beta-producing cells link innate and adaptive immunity. *J Exp Med* **192**:219, 2000
350. Ito T et al: Plasmacytoid dendritic cells prime IL-10-producing T regulatory cells by inducible costimulator ligand. *J Exp Med* **204**:105, 2007
351. de Heer HJ et al: Essential role of lung plasmacytoid dendritic cells in preventing asthmatic reactions to harmless inhaled antigen. *J Exp Med* **200**:89, 2004
352. Blanco P et al: Induction of dendritic cell differentiation by IFN-alpha in systemic lupus erythematosus. *Science* **294**:1540, 2001
353. Jego G et al: Plasmacytoid dendritic cells induce plasma cell differentiation through type I interferon and interleukin 6. *Immunity* **19**:225, 2003
354. Chaperot L et al: Virus or TLR agonists induce TRAIL-mediated cytotoxic activity of plasmacytoid dendritic cells. *J Immunol* **176**:248, 2006

355. Stary G et al: Plasmacytoid dendritic cells express TRAIL and induce CD4+ T-cell apoptosis in HIV-1 viremic patients. *Blood* **114**:3854, 2009
356. Riboldi E et al: Engagement of BDCA-2 blocks TRAIL-mediated cytotoxic activity of plasmacytoid dendritic cells. *Immunobiology* **214**:868, 2009
357. Piccioli D et al: Human plasmacytoid dendritic cells are unresponsive to bacterial stimulation and require a novel type of cooperation with myeloid dendritic cells for maturation. *Blood* **113**:4232, 2009
358. Novak N et al: Characterization of FcεpsilonRI-bearing CD123 blood dendritic cell antigen-2 plasmacytoid dendritic cells in atopic dermatitis. *J Allergy Clin Immunol* **114**:364, 2004
359. Fuchs A et al: Paradoxical inhibition of human natural interferon-producing cells by the activating receptor NKp44. *Blood* **106**:2076, 2005
360. Rissoan MC et al: Reciprocal control of T helper cell and dendritic cell differentiation. *Science* **283**:1183, 1999
361. Tel J et al: Human plasmacytoid dendritic cells phagocytose, process, and present exogenous particulate antigen. *J Immunol* **184**:4276, 2010
362. Aspord C et al: A novel cancer vaccine strategy based on HLA-A*0201 matched allogeneic plasmacytoid dendritic cells. *PLoS One* **5**:e10458, 2010