

# Chapter 123

## Atypical (Dysplastic) Melanocytic Nevi

James M. Grichnik & Margaret A. Tucker

### REFERENCES

1. Clark WH Jr et al: Origin of familial malignant melanoma from heritable melanocytic lesions. 'The B-K mole syndrome.' *Arch Dermatol* **114**:732, 1978 [PMID: 646394]
2. Lynch HT, Frichot BC III, Lynch JT: Familial atypical multiple mole-melanoma syndrome. *J Med Genet* **15**:352, 1978 [PMID: 739524]
3. Slade J et al: Atypical mole syndrome: Risk factor for cutaneous malignant melanoma and implications for management. *J Am Acad Dermatol* **32**:479, 1995 [PMID: 7868720]
4. Ackerman AB, Magana-Garcia M: Naming acquired melanocytic nevi: Unna's, Miescher's, Spitz's, Clark's. *Am J Dermatopathol* **12**:193, 1990 [PMID: 2184692]
5. NIH Consensus Development Panel on Early Melanoma: Diagnosis and treatment of early melanoma. *JAMA* **268**:1314, 1992
6. Naeyaert JM, Brochez L: Dysplastic nevi. *N Engl J Med* **349**:23, 2003
7. Shapiro M et al: Variability in nomenclature used for nevi with architectural disorder and cytologic atypia (microscopically dysplastic nevi) by dermatologists and dermatopathologists. *J Cutan Pathol* **31**:523, 2004 [PMID: 15268706]
8. Greene MH et al: Clinical recognition of acquired precursors of cutaneous malignant melanoma: A color atlas of the familial dysplastic nevus syndrome. *N Engl J Med* **312**:91, 1985 [PMID: 3964923]
9. Bergman W, Palan A, Went LN: Clinical and genetic studies in six Dutch kindreds with the dysplastic naevus syndrome. *Ann Hum Genet* **50**:249, 1986 [PMID: 3446012]
10. MacKie RM et al: CDKN2A germline mutations in UK patients with familial melanoma and multiple primary melanomas. *J Invest Derm* **111**:269, 1998 [PMID: 9699728]
11. Chaudru V et al: Influence of genes nevi, and sun sensitivity on melanoma risk in a family sample unselected by family history and in melanoma-prone families. *J Natl Cancer Inst* **96**:785, 2004 [PMID: 15150307]
12. Platz A et al: Screening of germline mutations in the CDKN2A and CDKN2B genes in Swedish families with hereditary cutaneous melanoma. *J Natl Cancer Inst* **89**:697, 1997 [PMID: 9168184]
13. Newton-Bishop JA et al: Family studies in melanoma: Identification of the atypical mole syndrome (AMS) phenotype. *Melanoma Res* **4**:199, 1994 [PMID: 7950355]
14. Ruiz A et al: CDKN2A mutations in Spanish cutaneous malignant melanoma families and patients with multiple and neoplasia. *J Med Genet* **36**:490, 1999 [PMID: 10874641]
15. Ghiorzo P et al: Characterization of Ligurian melanoma families and risk of occurrence of other neoplasia. *Int J Cancer* **83**:441, 1999 [PMID: 10508477]
16. Landi MT et al: Clinical characteristics of 20 Italian melanoma-prone families. *Arch Dermatol* **135**:1554, 1999 [PMID: 10606077]
17. Cooke KR et al: Dysplastic nevi in a population-based survey. *Cancer* **63**:1240, 1989 [PMID: 2917325]
18. Bataille V et al: The association between naevi and melanoma in populations with different levels of sun exposure: A joint case-control study of melanoma in the UK and Australia. *Br J Cancer* **77**:505, 1998 [PMID: 9472652]
19. Hara K, Nitta Y, Ikeya T: Dysplastic nevus syndrome among Japanese: A case study and review of the Japanese literature. *Am J Dermatopathol* **14**:24, 1992 [PMID: 1510212]
20. Goldstein AM et al: The inheritance pattern of dysplastic naevi in families of dysplastic naevus patients. *Melanoma Res* **3**:15, 1993 [PMID: 8471833]
21. Celebi JT et al: Evaluation of germline CDKN2A, ARF, CDK4, PTEN, and BRAF alterations in atypical mole syndrome. *Clin Experim Dermatol* **30**:68, 2004
22. James MR et al: BRAF polymorphisms and risk of melanocytic neoplasia. *J Invest Dermatol* **125**:1252, 2005 [PMID: 16354196]

23. Goldstein AM et al: Gene-covariate interaction between dysplastic nevi and the CDKN2A gene in American melanoma-prone families. *Cancer Epidemiol Biomarkers Prev* 9:889, 2000 [PMID: 11008905]
24. Falchi M et al: Genome-wide search for nevus density shows linkage to two melanoma loci on chromosome 9 and identifies a new QTL on 5q31 in an adult twin cohort. *Hum Molec Genet* 15:2975, 2006 [PMID: 16928783]
25. Zhu G et al: A genome-wide scan for naevus count: Linkage to CDKN2A and to other chromosomal regions. *Eur J Hum Genet* 15:94, 2007 [PMID: 17063143]
26. Falchi M et al: Genome-wide association study identifies variants at 9p21 and 22q13 associated with development of cutaneous nevi. *Nat Genet* 41:915, 2009
27. Bishop DT et al: Genome-wide association study identifies three loci associated with melanoma risk. *Nat Genet* 41:920, 2009
28. Pollack PM et al: High frequency of BRAF mutations in nevi. *Nat Genet* 33:19, 2003
29. Turner DJ et al: Detection of the BRAF V600E mutation in melanocytic lesions using the ligase detection reaction. *J Cutan Pathol* 32:334, 2005 [PMID: 1581117]
30. Eskandarpour M et al: Frequency of UV-inducible NRAS mutations in melanoma of patients with germline CDKN2A mutations. *J Natl Cancer Inst* 95:790, 2003 [PMID: 12783933]
31. Park WS et al: Allelic deletion at chromosome 9p21 (p16) and 17p13(p53) in microdissected sporadic dysplastic nevus. *Hum Pathol* 29:127, 1998 [PMID: 9490270]
32. Kopf AW et al: Relationship of nevocytic nevi to sun exposure in dysplastic nevus syndrome. *J Am Acad Dermatol* 12:656, 1985 [PMID: 3989027]
33. Kopf AW et al: Relationship of lumbosacral nevocytic nevi to sun exposure in dysplastic nevus syndrome. *Arch Dermatol* 122:1003, 1986 [PMID: 3740881]
34. Stierner U et al: Regional distribution of common and dysplastic nevi in relation to melanoma site and sun exposure. A case-control study. *Melanoma Res* 1:367, 1991
35. Tucker MA et al: A natural history of melanomas and dysplastic nevi: An atlas of lesions in melanoma-prone families. *Cancer* 94:3192, 2002 [PMID: 12115352]
36. Weinstock MA et al: Sunlight and dysplastic nevus risk: Results of a clinic-based case-control study. *Cancer* 767:1701, 1991
37. Titus-Ernstoff L et al: A relation between childhood sun exposure and dysplastic nevus syndrome among patients with nonfamilial melanoma. *Epidemiology* 2:210, 1991 [PMID: 2054404]
38. Garbe C et al: Associated factors in the prevalence of more than 50 common nevi, atypical melanocytic nevi, and actinic lentiginos: Multicenter case-control study of the Central Malignant Melanoma Registry of the German Dermatological Society. *J Invest Dermatol* 102:700, 1994 [PMID: 8176251]
39. Novakovic B et al: Melanocytic nevi, dysplastic nevi, and malignant melanoma in children from melanoma-prone families. *J Am Acad Dermatol* 33:631, 1995 [PMID: 7673498]
40. Tucker MA et al: Dysplastic nevi on the scalp of prepubertal children from melanoma-prone families. *J Pediatr* 103:65, 1983 [PMID: 6864397]
41. Halpern AC et al: A cohort study of melanoma in patients with dysplastic nevi. *J Invest Dermatol* 100:346S, 1993
42. Wang SQ et al: Detection of melanomas in patients followed up with total cutaneous examinations, total cutaneous photography, and dermoscopy. *J Am Acad Dermatol* 50:15, 2004 [PMID: 14699359]
43. Banky JP et al: Incidence of new and changed nevi and melanoma detected using baseline images and dermoscopy in patient at high risk for melanoma. *Arch Dermatol* 141:998, 2005 [PMID: 16103329]
44. Tucker MA et al: Clinically recognized dysplastic nevi: A central risk factor for cutaneous melanoma. *JAMA* 277:1439, 1997 [PMID: 9145715]
45. Elder DE, Clark WH Jr, Elenitsas R: The early and intermediate precursor lesions of tumor progression in the melanocytic system: Common acquired nevi and atypical (dysplastic) nevi. *Semin Diag Pathol* 10:18, 1993 [PMID: 8506414]
46. Hartge P et al: Recognition and classification of clinically dysplastic nevi from photographs: A study of interobserver variation. *Cancer Epidemiol Biomarkers Prev* 4:37, 1995 [PMID: 7894322]
47. Clemente C et al: Histopathologic diagnosis of dysplastic nevi: Concordance among pathologists convened by the World Health Organization Melanoma Programme. *Hum Pathol* 22:313, 1991 [PMID: 1741810]
48. deWit PE et al: Validity of the histopathological criteria used for diagnosing dysplastic naevi: An interobserver study by the pathology subgroups of the EORTC Malignant Melanoma Cooperative Group. *Er J Cancer* 29A:831, 1993

49. Weinstock MA et al: Reliability of the histopathologic diagnosis of melanocytic dysplasia. *Arch Dermatol* 133:953, 1997 [PMID: 9267239]
50. Lucas CR et al: Early melanoma detection: Non-uniform dermoscopic features and growth. *J Am Acad Dermatol* 48:663, 2003
51. Grob JJ, Bonerandi JJ: The 'ugly duckling' sign: Identification of the common characteristics of nevi in an individual as a basis for melanoma screening. *Arch Dermatol* 134:103, 1998
52. Fears TR et al: Identifying individuals at high risk of melanoma: A practical predictor of absolute risk. *J Clin Oncol* 24:3590, 2006 [PMID: 16728488]
53. Chaudru V et al: Melanocortin-1 receptor (MC1R) gene variants and dysplastic nevi modify penetrance of CDKN2A mutations in French melanoma-prone pedigrees. *Cancer Epidemiol Biomarkers Prev* 14:2384, 2005 [PMID: 16214921]
54. Elder DE: Precursors to melanoma and their mimics: Nevi of special sites. *Modern Pathol* 19:S4, 2006
55. Clark WH Jr et al: Atypical melanocytic nevi of the genital type with a discussion of reciprocal parenchymal-stromal interactions in the biology of neoplasia. *Hum Pathol* 29:S1, 1998
56. Sagebiel RW: The dysplastic melanocytic nevus. *J Am Acad Dermatol* 20:496, 1989 [PMID: 2645327]
57. Barnhill RL: Melanocytic nevi and tumor progression: Perspectives concerning histomorphology, melanoma risk, and molecular genetics. *Dermatology* 187:86, 1993 [PMID: 8358111]
58. Murphy GF, Mihm MC: Recognition and evaluation of cytological dysplasia in acquired melanocytic nevi. *Hum Pathol* 30:506, 1999 [PMID: 10333218]
59. Pehamberger H et al: In vivo epiluminescence microscopy: Improvement of early diagnosis of melanoma. *J Invest Dermatol* 100:356S, 1993
60. Nachbar F et al: The ABCD rule of dermatoscopy. High prospective value in the diagnosis of doubtful melanocytic skin lesions. *J Am Acad Dermatol* 30:551, 1994
61. Menzies SW et al: Frequency and morphologic characteristics of invasive melanomas lacking specific surface microscopic features. *Arch Dermatol* 132:1178, 1996
62. Argenziano G et al: Epiluminescence microscopy for the diagnosis of doubtful melanocytic skin lesions. Comparison of the ABCD rule of dermatoscopy and a new 7-point checklist based on pattern analysis. *Arch Dermatol* 134:1563, 1998
63. Henning JS et al: The CASH (color, architecture, symmetry, and homogeneity) algorithm for dermoscopy. *J Am Acad Dermatol*. 56:45, 2007
64. Vestergaard ME et al: Dermoscopy compared with naked eye examination for the diagnosis of primary melanoma: A meta-analysis of studies performed in a clinical setting. *Br J Dermatol*. 159:669, 2008
65. Kittler H, Binder M: Risks and benefits of sequential imaging of melanocytic skin lesion in patients with multiple atypical nevi. *Arch Dermatol* 137:1590, 2001 [PMID: 11735709]
66. Menzies SW et al: Short-term digital surface microscopic monitoring of atypical or changing melanocytic lesions. *Arch Dermatol* 137:1583, 2001 [PMID: 11735708]
67. Kopf AW et al: Photographs are useful for detection of malignant melanomas in patients who have dysplastic nevi. *J Am Acad Dermatol* 19:1132, 1988
68. Masri GD et al: Screening and surveillance of patients at high risk for malignant melanoma result in detection of earlier disease. *J Am Acad Dermatol* 22:1042, 1990 [PMID: 2370329]
69. MacKie RM, McHenry P, Hole D: Accelerated detection with prospective surveillance for cutaneous malignant melanoma in high-risk groups. *Lancet* 341:1618, 1993 [PMID: 8099990]
70. Schneider JS, Moore DH, Sagebiel RW: Risk factors for melanoma incidence in prospective follow-up: The importance of atypical (dysplastic) nevi. *Arch Dermatol* 130:1002, 1994 [PMID: 8053695]
71. Kelly JW et al: A high incidence of melanoma found in patients with multiple dysplastic naevi by photographic surveillance. *MJA* 167:191, 1997 [PMID: 9293264]
72. Busam KJ, Marghoob AA, Halpern A: Melanoma diagnosis by confocal microscopy: Promise and pitfalls. *J Invest Dermatol* 125:vii, 2005
73. Pellacani G, Cesarino AM, Seidenari S: Reflectance-mode confocal microscopy for the in vivo characterization of pagetoid melanocytosis in melanomas and nevi. *J Invest Dermatol* 125:532, 2005 [PMID: 16117795]
74. Friedman RJ et al: The diagnostic performance of expert dermoscopists vs a computer-vision system on small-diameter melanomas. *Arch Dermatol* 144:476, 2008
75. Thomas MW, Grichnik JM, Izatt JA: Three-dimensional images and vessel rendering using optical coherence tomography. *Arch Dermatol*. 143:1468, 2007

76. Guitera P et al: Melanoma histological Breslow thickness predicted by 75-MHz ultrasonography. *Br J Dermatol*. **159**:364, 2008
77. Zhao J et al: Real-time Raman spectroscopy for non-invasive skin cancer detection—Preliminary results. *Conf Proc IEEE Eng Med Biol Soc* **2008**:3107, 2008
78. Han X et al: Near-infrared autofluorescence imaging of cutaneous melanins and human skin in vivo. *J Biomed Opt*. **14**:024017, 2009
79. Tucker MA, Goldstein AM: Melanoma etiology: Where are we? *Oncogene* **22**:3042, 2003 [PMID: 12789279]
80. Bauer J, Garbe C: Acquired melanocytic nevi as risk factor for melanoma development. A comprehensive review of epidemiologic data. *Pigment Cell Res* **16**:297, 2003 [PMID: 12753404]
81. Gandini S et al: Meta-analysis of risk factors for cutaneous melanoma: I. Common and atypical naevi. *Eur J Cancer* **41**:28, 2005 [PMID: 15617989]
82. Freedman DM, Miller BA, Tucker MA: New malignancies following melanoma of the skin, eye melanoma, and non-melanoma eye cancer. In: *New Malignancies Among Cancer Survivors: SEER Cancer Registries, 1973–2000*, edited by RE Curtis et al Bethesda, National Cancer Institute, NIH Publication No. 05-5302, 2006, p. 339
83. Ferrone CR et al: Clinicopathological features of and risk factors for multiple primary melanomas. *JAMA* **294**:1647, 2005 [PMID: 16204664]
84. Gandini S et al: Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer* **41**:45, 2005
85. Lee TK, Rivers JK, Gallagher RP: Site-specific protective effect of broad-spectrum sunscreen on nevus development among white schoolchildren in a randomized trial. *J Am Acad Dermatol* **52**:786, 2005 [PMID: 15858467]